

1. **Scaling of the dynamical equations** (3 points)

We work in the rotating frame of reference of the Earth. The equation can be scaled by a length-scale  $L$ , determined by the geometry of the flow, and by a characteristic velocity  $U$ . We can estimate the relative contributions in units of  $m/s^2$  in the horizontal momentum equations:

$$\underbrace{\frac{\partial \mathbf{v}}{\partial t}}_{U/T \sim 10^{-8}} + \underbrace{\mathbf{v} \cdot \nabla \mathbf{v}}_{U^2/L \sim 10^{-8}} = \underbrace{-\frac{1}{\rho} \nabla p}_{\delta P / (\rho L) \sim 10^{-5}} + \underbrace{2\boldsymbol{\Omega} \times \mathbf{v}}_{f_0 U \sim 10^{-5}} + \underbrace{fric}_{\nu U / H^2 \sim 10^{-13}} \quad (1)$$

where  $fric$  denotes the contributions of friction due to eddy stress divergence (usually  $\sim \nu \nabla^2 \mathbf{v}$ ). Typical values are given in Table 1. The values have been taken for the ocean.

a) Please repeat the estimate for the atmosphere using Table 1.

b) The Rossby number  $Ro$  is the ratio of inertial (the left hand side in (1)) to Coriolis (second term on the right hand side in (1)) terms

$$Ro = \frac{(U^2/L)}{(fU)} = \frac{U}{fL} \quad (2)$$

$Ro$  is small when the flow is in a so-called geostrophic balance. Please calculate  $Ro$  for the atmosphere and ocean using Table 1.

	Quantity	Atmosphere	Ocean
horizontal velocity	$U$	$10 \text{ ms}^{-1}$	$10^{-1} \text{ ms}^{-1}$
horizontal length	$L$	$10^6 \text{ m}$	$10^6 \text{ m}$
horizontal Pressure changes	$\delta P$ (horizontal)	$10^3 \text{ Pa}$	$10^4 \text{ Pa}$
time scale	$T$	$10^5 \text{ s}$	$10^7 \text{ s}$
Coriolis parameter at $45^\circ\text{N}$	$f_0 = 2\Omega \sin \varphi_0$	$10^{-4} \text{ s}^{-1}$	$10^{-4} \text{ s}^{-1}$
density	$\rho$	$1 \text{ kgm}^{-3}$	$10^3 \text{ kgm}^{-3}$
viscosity (turbulent)	$\nu$	$10^{-5} \text{ kgm}^{-3}$	$10^{-6} \text{ kgm}^{-3}$

Table 1: Table shows the typical scales in the atmosphere and ocean system.

2. **Concept of dynamic similarity** (4 points)

For the case of an incompressible flow, assuming the temperature effects are negligible and external forces are neglected, the Navier-Stokes equations consist of conservation of mass

$$\nabla \cdot \mathbf{u} = 0 \quad (3)$$

and conservation of momentum

$$\partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\frac{1}{\rho_0} \nabla p + \nu \nabla^2 \mathbf{u} \quad (4)$$

where  $\mathbf{u}$  is the velocity vector and  $p$  is the pressure,  $\nu$  denotes the kinematic viscosity.

a) Show: The equations (3,4) can be made dimensionless by a length-scale  $L$ , determined by the geometry of the flow, and by a characteristic velocity  $U$ . For example:  $u = U \cdot u_d$ .

Note: the units of  $[\rho_0] = kg/m^3$ ,  $[p] = kg/(ms^2)$ , and  $[p]/[\rho_0] = m^2/s^2$ . Therefore the pressure gradient term in (4) has the scaling  $U^2/L$ .

b) Show: The scalings vanish completely in front of the terms except for the  $\nabla^2 \mathbf{u}_d$ -term! The dimensionless parameter is the Reynolds number and the only parameter left!

*Remark: For large Reynolds numbers, the flow is turbulent. In most practical flows  $Re$  is rather large ( $10^4 - 10^8$ ), large enough for the flow to be turbulent.*

3. **Advection** (3 points)

A ship is steaming northward at a rate of 10 km/h. The surface pressure increases toward the northwest at a rate of 5 Pa/km. What is the pressure tendency recorded at a nearby island station if the pressure aboard the ship decreases at a rate of 100Pa/3h?

Notes on submission form of the exercises: Working in study groups is encouraged, but each student is responsible for his/her own solution. The answers to the questions can be send until the due date to Akil Hossain (akil.hossain@awi.de).