

1. **Wind-driven ocean circulation** (6 points)

The **Sverdrup transport** V for the depth-integrated flow is calculated by

$$\rho_0 \beta V = \frac{\partial}{\partial x} \tau_y - \frac{\partial}{\partial y} \tau_x \quad (1)$$

where τ_x and τ_y are the components of the wind stress.

The **Ekman transports** V_E, U_E describe the dynamics in the upper mixed layer:

$$f V_E = -\tau_x / \rho_0, \quad f U_E = \tau_y / \rho_0 \quad (2)$$

where $U_E = \int_{-E}^0 u dz$ and $V_E = \int_{-E}^0 v dz$ are the depth-integrated velocities in the thin friction-dominated Ekman layer at the sea surface.

Ekman vertical velocity w_E : Using the continuity equation, the divergence of the Ekman transports leads to a vertical velocity w_E at the bottom of the Ekman layer:

$$w_E = - \int_{-E}^0 \frac{\partial w}{\partial z} dz = \frac{\partial}{\partial x} U_E + \frac{\partial}{\partial y} V_E = \frac{\partial}{\partial x} \left(\frac{\tau_y}{\rho_0 f} \right) - \frac{\partial}{\partial y} \left(\frac{\tau_x}{\rho_0 f} \right) \quad (3)$$

a) Assume that the windstress is only zonal with

$$\tau_x = -\tau_0 \cos(\pi y / B) \quad \text{for an ocean basin with } 0 < x < L, \ 0 < y < B. \quad (4)$$

Calculate the Sverdrup transport, Ekman transports, and Ekman pumping velocity for this special case.

b) Make a schematic diagram of the windstress, Sverdrup transport, Ekman transports, and Ekman pumping velocity.

c) Using a), at what latitudes y are $|V|$ and $|V_E|$ maximum? Calculate their magnitudes. Take constant $f = 10^{-4} \text{ s}^{-1}$ and $\beta = 1.8 \cdot 10^{-11} \text{ m}^{-1} \text{ s}^{-1}$ and $B = 5000 \text{ km}$, $\tau_0 / \rho_0 = 10^{-4} \text{ m}^2 \text{ s}^{-2}$.

d) Using the values in b), calculate the maximum of w_E for constant f .

2. **Rossby wave formula (long waves in the westerlies)** (4 points)

a) Assume a mean flow with constant zonal velocity $u = U = \text{const} > 0$ and a varying north-south component $v = v(x, t)$ which gives the total motion a wave-like form. Furthermore, $h = \text{const}$.

Write down the vorticity equation for this specific flow! Remember that the vorticity equation is

$$\frac{D}{Dt} \left(\frac{\zeta + f}{h} \right) = 0 \quad (5)$$

b) Use a) and the ansatz

$$v(x, t) = A \cos[(kx - \omega t)] \quad (6)$$

to determine the dispersion relation $\omega(k)$, the group velocity $\frac{\partial \omega}{\partial k}$, and the phase velocity $c = \omega/k$.

3. Conservation of potential vorticity: (2 points)

An air column at 53°N with $\zeta = 0$ initially stretches from the surface to a fixed tropopause at 10 km height. If the air column moves until it is over a mountain barrier of 2 km height at 30°N , what is its absolute vorticity and relative vorticity as it passes the mountain top?

Assume: $\sin 53^\circ = 0.8$; $\sin 30^\circ = 0.5$

The angular velocity of the Earth $\Omega = 2\pi/(1 \text{ day})$.

Potential vorticity: $(\zeta + f)/h$

Notes on submission form of the exercises: *Working in study groups is encouraged, but each student is responsible for his/her own solution. The answers to the questions can be send until the due date (12:00) to Aparna Prasannakumar (apa_pra@uni-bremen.de) and Yifan Ma (yifan.ma@awi.de).*