

Mathematics and Climate Change

Gerrit Lohmann

Contents

Introduction	2
Climate: A Fluid Dynamical System	6
Mathematical Equations	6
Nondimensional Parameters: The Reynolds Number	9
Convection in the Rayleigh-Bénard System	11
Reduction of Dimensions and the Lorenz System	13
Scaling in the Climate System	15
Projection Methods: Coarse Graining and Stable Manifold Theory	16
Brownian Motion, Weather, and Climate	18
Climate Variability and Sensitivity	19
Non-normal Growth of the Climate System	21
Predictability	24
Boltzmann Dynamics	26
Conclusions	29
Cross-References	30
References	30

Abstract

Climate change is one of the most pressing scientific challenges of our times, with transformations which are already becoming present in many areas of the world. The demand (from the stakeholders) for clear answers under a wide range of future scenarios has to be addressed (by the scientific community) using our rapidly evolving knowledge of the weather and climate system. Mathematics is

G. Lohmann (🖂)

Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany

University of Bremen, Bremen, Germany

© Springer Nature Switzerland AG 2021

B. Sriraman (ed.), Handbook of the Mathematics of the Arts and Sciences, https://doi.org/10.1007/978-3-319-70658-0_145-1

e-mail: Gerrit.Lohmann@awi.de

one of the essential pillars at the foundation of this knowledge, as it allows us to quantify and predict the effects we observe in nature. In this chapter, we illustrate some crucial mathematical techniques and theoretical approaches, in the context of their application to the climate system. The concepts of critical parameters, dimension reduction, and stochasticity are explored in detail.

Keywords

Climate models · Fluid dynamics · Dynamical theory · Spatiotemporal scales · Coarse graining · Stability · Predictability

Introduction

Mathematics can be described in large parts on (unprovable, but very well founded) axioms by purely logical steps. Physics owes, since the beginning of modern times, its great successes due to experiments (can be repeated at any time with limitation to measurable data) with mathematical theories and models. Strong abstraction, e.g., in free fall, is unavoidable and at the same time greatly reduces the "overall reality." Climate science is rather new subject, describing the nature of its components and quantities like temperatures and currents. Its great success is due to a proper combination of observations, its theoretical foundations in fluid dynamics, and the statistical analysis of data. Unlike in physics, there is no lab to repeat measurements; instead, we have just one realization of the climate trajectory. Until now it is unknown on whether the gradual or the catastrophic case is more likely.

Figure 1 shows the Northern Hemisphere temperature evolution of the last 150 years. The last 150 years are quite often called the "instrumental period" since the spatial coverage and the quality of the data is high compared to earlier periods. The current and future climate is subject to significant change and fluctuations; a large



Fig. 1 Northern Hemisphere near-surface temperature anomaly [K] based on HadCRUT4 (Morice et al., 2012)

part is due to the increasing human influence on the climate system. The extent and the rate of this change are controversial, however. It is therefore necessary to improve the understanding of natural climate variability and trends by searching for their causes at time different scales, i.e., the multidecadal component in Fig. 1. A major challenge is furthermore to understand the dynamics and potential thresholds of rapid climate changes. The analysis of the current status, of the past, of driving mechanisms and feedbacks provides a suitable framework to study conditions which are expected to develop in the future.

A comprehensive modeling strategy designed to address abrupt climate change includes vigorous use of a hierarchy of models, from theory and conceptual models with only a few degrees of freedom through models of intermediate complexity, to high-resolution models of components of the climate system, to fully coupled Earth system models. The simpler models are well-suited for use in developing new hypotheses for abrupt climate change. Model-data comparisons are needed to assess the quality of model predictions. It is important to note that the multiple long integrations of enhanced, fully coupled Earth system models required for this research are not possible with the computer resources available today, and thus, these resources are currently enhanced. Since Earth system models have to simplify the system and rely on parameterizations of unresolved processes using present data, paleoclimate records provide a unique tool to validate models for conditions which are different from our present one. Suitable model-data analyses provide therefore a proper basis to estimate and possibly reduce uncertainties of future climate change projections (Lohmann et al., 2020). Furthermore, the model scenarios in conjunction with the long-term data can be used to examine mechanisms for the statistics of regional climate extremes under different boundary conditions. Mathematical tools are numerics of partial differential equations, and conceptual approaches of fluid mechanics are described in this chapter.

In the entire climate system, different scales play an important role (Fig. 2). These are the characteristic orders of magnitude in space and time that a system possesses or is superimposed on the system in order to record or observe it. Climate has a spatial and temporal dimension, which fluctuate in a wide range of spatial and temporal scales. Spatial scales vary from local to regional to continental. Time scales vary from seasonal to geological. The spatiotemporal dimension of complex phenomena are defined by their typical spatial extension (e.g., the diameter of a high-pressure area), which is linked to a structure whose magnitudes can be specified as spatial scales. The time scale of an atmospheric process is the order of magnitude of its lifetime. In addition, it is also possible to specify the spatial and temporal resolution with which a system is to be viewed. A distinction is made between the scale in space and time that an atmospheric process or an atmospheric system has and the scale of the observation. If the spatial and temporal scales of observation are to make it possible to capture the system in good resolution, they must be significantly smaller than those describing the overall system. The word "scale" in this context means order of magnitude or scale. The spectrum of atmospheric space and time scales covers many orders of magnitude. A rough classification, which is common in meteorology, is based on the horizontal space



Fig. 2 Schematic diagram of the spatio-temporal scales considered. *DO*: Dansgaard-Oeschger, *H*: Heinrich events; *AMO*: Atlantic Multidecadal Oscillation; *PDO*: Pacific Decadal Oscillation; *ENSO*: El Niño-Southern Oscillation. The annual and astronomical cycles are externally driven and have quasi-global impact. The dashed line shows a schematic power spectrum with more variability on long time scales

scale L. Basically, climate represents a space-time continuum, so that fixed scale limits do not occur in the real atmosphere. Rather, more or less narrow transition ranges between the scales are the rule. Larger and smaller systems influence each other so that the transitions between differently scaled weather phenomena are smooth and the approach is usually based on the question. In order to illustrate the scaling in the climate system, the procedure of non-dimensional parameters are introduced.

Models of planetary motion based on Newton's models of gravity and motion were astonishingly successful, and this had a profound impact on the way in which people viewed mathematical models and in the way that we still view them. The eighteenth-century French scientist/mathematician Laplace, extended the basic Newtonian model, given above, to model the motion of all the planets in the solar system, including their influences on each other. This model accounted for the motions of the planets as perfectly as they could be measured (including all the small deviations from elliptical orbits caused by the planets gravitational effects on each other). This seemed to be such a triumph that the belief grew up that everything in the universe could be described by such models and that in principle the future could be predicted perfectly given such models and accurate measurements of the state of the universe now. Perhaps not surprisingly, Laplace was a vigorous promoter of this idea, which gained the name determinism.

These ideas lent their name to the concept of a deterministic model – that is, a model which if solved from identical starting conditions always has the same solution. For much of the nineteenth-century, determinism reined and resulted in a great deal of heart searching about free will and the like. In the twentieth century, three areas of science comprehensively overturned the idea that everything could be described by deterministic models. These were quantum mechanics and chaos theory. The later had its origin in meteorology and fluid dynamics.

In the 1960s, Edward Lorenz, a mathematician and meteorologist, showed that there are natural limits to the predictability of a nonlinear system, such as atmospheric circulation. He discovered and described the chaotic behavior of large-scale motion patterns in the atmosphere and showed that despite the determinacy of the system, i.e., that although the partial differential equations could be calculated at any time, the system itself loses its predictability after a relatively short time. Even the smallest changes in the initial conditions caused different final states after a few iteration steps (calculation steps). A predictability of the system is therefore limited in time and the nonlinearity is responsible for the finite predictability of atmospheric flow patterns. This insight became known under the technical term "butterfly effect".

Climate physicists are constantly trying to learn from our observations how to examine causes and effects in such a way that the observations are captured. We write down our gained insights as mathematically formulated laws of nature. We formulate equations of motion in the form of differential equations. Their solutions provide information about what will happen to a given state at a given time at a later time. In this sense, the equations of motion as such provide a causal description par excellence - all equations of motion. Sometimes, however, there seem to be annoying difficulties with causal chains or with predictability, for example, when the destructive path of a hurricane is poorly predicted, or in extreme weather events. We are even more aware of the problems in long-term predictions, such as climate change. So is causality failing here? Of course we do not think so, otherwise we would not be looking for causes. It then also made no sense to derive political decisions from insights into climate evolution. And although word has got around that quantum mechanics is not a causal physics in the classical sense, we will hardly want to blame quantum mechanics for the cases of lack of predictability or undetectable cause-effect relationships. Here, the intersection of mathematical modeling and questions within climate change sciences are elaborated.

A systematic description of the mathematics and climate is given (Fig. 3). A general question within the micro-macro dynamic is that of integration between different levels. Two distinctly different levels emerge with different rules governing each, but they then need to be reconciled in some way to create an overall functioning system. Physical, chemical, biological, economic, social, and cultural systems all exhibit this micro-macro dynamic and how the system comes to reconcile it forms a primary determinate in its identity and overall structure. This multidimensional nature to a system that results in the micro-macro dynamic is a product of synthesis and emergence. An approach is coarse graining and projection



Validation of climate scenarios on different time scales by using observations and paleoclimates

Fig. 3 Systematic description of climate and mathematics. Changing the description of the dynamics: from the micro- to macroscales. This is a common problem since we are not able to describe the systems on all temporal and spatial scales

where the underlying dynamics is projected onto the macroscopic dynamics; the other is the statistical physics theory of nonequilibrium statistical mechanics. The Boltzmann equation, coarse graining, and the Brownian motion are the approaches to understand the dynamics on different scales.

Climate: A Fluid Dynamical System

For present climate state, we are able to directly measure all involved quantities. From measurements we can draw conclusions about physical, chemical, and biological relationships between the variables. Our understanding about the involved processes is far from complete, but nevertheless we derive equations that describe and predict the observed phenomena (Fig. 4).

Mathematical Equations

Our starting point is a mathematical model for the system of interest. In physics a model typically describes the state variables, plus fundamental laws and equations of state. These variables evolve in space and time. For the ocean, fundamental equations are formulated:



Fig. 4 Schematic view on the climate system. Global climate is a result of the complex interactions between the atmosphere, cryosphere (ice), hydrosphere (oceans), lithosphere (land), and biosphere (life), fueled by the nonuniform spatial distribution of incoming solar radiation (Peixoto and Oort, 1992). We know from climate reconstructions using recorders such as ice cores, ocean and lake sediment cores, tree rings, corals, cave deposits, and groundwater that the Earth's climate has seen major changes over its history. An analysis of the temperature variations patched together from all these data reveals that climate change occurs in cycles with characteristic periods, for example, 200 million, 100 thousand, or 4–7 years. For some of these cycles, particular mechanisms can be identified, for example, forcing by changes in the Earth's orbital parameters or internal oscillations of the coupled ocean-atmosphere system. However, major uncertainties remain in our understanding of the interplay of the components of the climate system

- State variables: Velocity (in each of three directions), pressure, temperature, salinity, and density
- Fundamental laws: Conservation of momentum, conservation of mass, conservation of temperature, and salinity
- Equations of state: Relationship of density to temperature, salinity, and pressure and perhaps also a model for the formation of sea ice

The state variables are expressed as a continuum in space and time and the fundamental laws as partial differential equations. If the atmosphere is becoming too thin in the upper levels, a more molecular, statistical description is appropriate. Even at this stage, though, simplifications may be made. For example, it is common to treat seawater as incompressible. Furthermore, equations of state are often specified by empirical relationships or laboratory experiments.

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{u}) = 0 \tag{1}$$

or, using the substantive derivative:

$$\frac{D\rho}{Dt} + \rho(\nabla \cdot \mathbf{u}) = 0.$$
⁽²⁾

A simplification of the resulting flow equations is obtained when considering an incompressible flow of a Newtonian fluid. The assumption of incompressibility rules out the possibility of sound or shock waves to occur, so this simplification is invalid if these phenomena are important. The incompressible flow assumption typically holds well even when dealing with a "compressible" fluid – such as air at room temperature – at low Mach numbers (even when flowing up to about Mach 0.3).

The dynamics of flow are based on the Navier-Stokes equations. This is a statement of the conservation of momentum in a fluid and it is an application of Newton's second law to a continuum; in fact this equation is applicable to any non-relativistic continuum and is known as the Cauchy momentum equation (e.g., Landau and Lifshitz 1959). Taking this into account and assuming constant viscosity, the Navier-Stokes equations will read, in vector form:



Note that only the advection terms are nonlinear for incompressible Newtonian flow. This acceleration is an acceleration caused by a (possibly steady) change in velocity over position, for example, the speeding up of fluid entering a converging nozzle. Though individual fluid particles are being accelerated and thus are under unsteady motion, the flow field (a velocity distribution) will not necessarily be time dependent.

The vector field **F** represents "other" (body force) forces. Typically this is only gravity but may include other fields (such as electromagnetic). In a non-inertial coordinate system, other "forces" such as that associated with rotating coordinates may be inserted. We note that the Coriolis force will be one of the main contributions in the rotating Earth system. Often, these forces may be represented as the gradient of some scalar quantity. Gravity in the z direction, for example, is the gradient of $-\rho gz$. Since pressure shows up only as a gradient, this implies that solving a problem without any such body force can be mended to include the body force by modifying pressure.

If temperature effects are also neglected, the only "other" equation (apart from initial/boundary conditions) needed is the mass continuity equation. Under the

incompressible assumption, density is a constant and it follows that the equation will simplify to:

$$\nabla \cdot \mathbf{u} = 0. \tag{4}$$

This is more specifically a statement of the conservation of volume (see divergence). These equations are commonly used in three coordinates systems: Cartesian, cylindrical, and spherical. While the Cartesian equations seem to follow directly from the vector equation above, the vector form of the Navier-Stokes equation involves some tensor calculus which means that writing it in other coordinate systems is not as simple as doing so for scalar equations (such as the heat equation).

Taking the curl of the Navier-Stokes equation results in the elimination of pressure. This is especially easy to see if two-dimensional Cartesian flow is assumed (w = 0 and no dependence of anything on z), where the equations reduce to:

$$D_t\left(\nabla^2\psi\right) = \nu\nabla^4\psi\tag{5}$$

where ∇^4 is the (2D) biharmonic operator and ν is the kinematic viscosity $\nu = \frac{\mu}{\rho}$. This single equation together with appropriate boundary conditions describes 2D fluid flow, taking only kinematic viscosity as a parameter. Note that the equation for creeping flow results when the left side is assumed zero. In axisymmetric flow another stream function formulation, called the Stokes stream function, can be used to describe the velocity components of an incompressible flow with one scalar function. The concept of taking the curl of the flow will become very important in climate dynamics (vorticity dynamics). The term $\zeta = \nabla^2 \psi$ is called relative vorticity, and the term $f = 2\Omega \sin \varphi$ is due to the rotating Earth (Ω is the radiation, φ the latitude). The dynamics can be described by the barotropic vorticity equation as

$$D_t(\zeta + f) = \nu \nabla^2 \zeta \tag{6}$$

which is heavily used in climate research.

Nondimensional Parameters: The Reynolds Number

In climate, we are interested in the critical parameters of the system. For the case of an incompressible flow in the Navier-Stokes equations, assuming the temperature effects are negligible and external forces are neglected, the equations consist of conservation of mass

$$\nabla \cdot \mathbf{u} = 0 \tag{7}$$

and momentum

$$\partial_t \mathbf{u} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\frac{1}{\rho_0} \nabla p + \nu \nabla^2 \mathbf{u}$$
(8)

where **u** is the velocity vector and p is the pressure and ν denotes the kinematic viscosity. The equations can be made dimensionless by a length scale L, determined by the geometry of the flow, and by a characteristic velocity U. For inter-comparison of analytical solutions, numerical results, and experimental measurements, it is useful to report the results in a dimensionless system. This is justified by the important concept of dynamic similarity (Buckingham, 1914). The main goal for using this system is to replace physical or numerical parameters with some dimensionless numbers, which completely determine the dynamical behavior of the system.

The procedure for converting to this system first implies, first of all, the selection of some representative values for the physical quantities involved in the original equations (in the physical system). For our current problem, we need to provide representative values for velocity (U), time (T), and distances (L). From these, we can derive scaling parameters for the time derivatives and spatial gradients also. Using these values, the values in the dimensionlesssystem (written with subscript d) can be defined:

$$u = U \cdot u_d \tag{9}$$

$$t = T \cdot t_d \tag{10}$$

$$x = L \cdot x_d \tag{11}$$

with U = L/T. From these scalings, we can also derive

$$\partial_t = \frac{\partial}{\partial t} = \frac{1}{T} \cdot \frac{\partial}{\partial t_d} \tag{12}$$

$$\partial_x = \frac{\partial}{\partial x} = \frac{1}{L} \cdot \frac{\partial}{\partial x_d} \tag{13}$$

Note furthermore the units of $[\rho_0] = kg/m^3$, $[p] = kg/(ms^2)$, and $[p]/[\rho_0] = m^2/s^2$. Therefore the pressure gradient term has the scaling U^2/L . Furthermore, divide the momentum equation by U^2/L and the scalings vanish completely in front of the terms except for the $\nabla_d^2 \mathbf{u}_d$ -term:

$$\nabla_d \cdot \mathbf{u_d} = 0 \tag{14}$$

and conservation of momentum

$$\frac{\partial}{\partial t_d} \mathbf{u_d} + (\mathbf{u_d} \cdot \nabla_d) \mathbf{u_d} = -\nabla_d p_d + \frac{1}{Re} \nabla_d^2 \mathbf{u_d}$$
(15)

The dimensionless parameter Re = UL/v is the Reynolds number and the only parameter left.

For large Reynolds numbers, the flow is turbulent. In most practical flows Re is rather large $(10^4 - 10^8)$, large enough for the flow to be turbulent. A large Reynolds number allows the flow to develop steep gradients locally. The typical length scale corresponding to these steep gradients can become so small that viscosity is not negligible. So the dissipation takes place at small scales. In this way different length scales are present in a turbulent flow, which range from L to the Kolmogorov length scale. This length scale is the typical length of the smallest eddy present in a turbulent flow. In the climate system, this dissipation by turbulence is modeled via eddy terms. To evaluate the critical parameters and scales, we implicitly assume such procedure. A classical example is provided in the next section.

Convection in the Rayleigh-Bénard System

A system of three ordinary differential equations is introduced whose solutions afford the simplest example of deterministic flow that we are aware of. The system is a simplification of the one derived by Saltzman (1962), to study finite-amplitude convection.

Consider the Rayleigh-Bénard circulation (Fig. 5). Rayleigh (1916) studied the flow occurring in a layer of fluid of uniform circulation depth H, when the temperature difference between the upper and lower surfaces is maintained at a constant value ΔT .

$$T(x, y, z = H) = T_0$$

 $T(x, y, z = 0) = T_0 + \Delta T$ (16)

The Boussinesq approximation is used, which results in a buoyancy force term which couples the thermal and fluid velocity fields. Therefore

$$\rho \approx \rho_0 = \text{const.}$$
(17)



Fig. 5 Geometry of the Rayleigh-Bénard system (see text for details)

except in the buoyancy term, where:

$$\rho = \rho_0 (1 - \alpha (T - T_0)) \text{ with } \alpha > 0.$$
 (18)

 ρ_0 is the fluid density in the reference state. This assumption reflects a common feature of geophysical flows, where the density fluctuations caused by temperature variations are small, yet they are the ones driving the overall flow. We have the following relations. Furthermore, we assume that the density depends linearly on temperature *T*.

This system possesses a steady-state solution in which there is no motion, and the temperature varies linearly with depth:

$$u = w = 0$$

$$T_{eq} = T_0 + \left(1 - \frac{z}{H}\right) \Delta T$$
(19)

When this solution becomes unstable, convection should develop.

In the case where all motions are parallel to the x - z-plane, and no variations in the direction of the y-axis occur, the governing equations may be written (see Saltzman 1962) as:

$$D_t u = -\frac{1}{\rho_0} \partial_x p + v \nabla^2 u \tag{20}$$

$$D_t w = -\frac{1}{\rho_0} \partial_z p + \nu \nabla^2 w + g(1 - \alpha (T - T_0))$$
(21)

$$D_t T = \kappa \nabla^2 T \tag{22}$$

$$\partial_x u + \partial_z w = 0 \tag{23}$$

where w and u are the vertical and horizontal components of the velocity. Furthermore, $v = \eta/\rho_0$, $\kappa = \lambda/(\rho_0 C_v)$ the momentum diffusivity (kinematic viscosity) and thermal diffusivity, respectively.

Now, the pressure is eliminated to derive the vorticity equation $D_t (\nabla^2 \psi) = \nu \nabla^4 \psi$. Here, it is useful to define the stream function Ψ for the two-dimensional motion, i.e.,

$$\frac{\partial \Psi}{\partial x} = w \tag{24}$$

$$\frac{\partial \Psi}{\partial z} = -u. \tag{25}$$

$$\frac{\partial}{\partial x}(21) - \frac{\partial}{\partial z}(20) = \frac{\partial}{\partial x}D_tw - \frac{\partial}{\partial z}D_tu = D_t\frac{\partial w}{\partial x} - D_t\frac{\partial u}{\partial z}$$
(26)

$$==D_t \frac{\partial^2 \Psi}{\partial x^2} - D_t \frac{\partial^2 \Psi}{\partial z^2} = D_t \nabla^2 \Psi.$$
 (27)

Furthermore, one can introduce the function Θ as the departure of temperature from that occurring in the state of no convection (19):

$$T = T_{eq} + \Theta \tag{28}$$

In the temperature term in $\frac{\partial}{\partial x}$ (21) on the right-hand side:

$$\frac{\partial}{\partial x}g(1-\alpha(T_{eq}+\Theta-T_0)) = -g\alpha\frac{\partial}{\partial x}\Theta$$

The left-hand side of (22) reads

$$D_t T = D_t T_{eq} + D_t \Theta = w \cdot \frac{-\Delta T}{H} + D_t \Theta = -\frac{\Delta T}{H} \frac{\partial \Psi}{\partial x} + D_t \Theta$$

Then, the dynamics can be formulated as

$$D_t\left(\nabla^2\Psi\right) = \nu\nabla^4\Psi - g\alpha\frac{\partial\Theta}{\partial x} \tag{29}$$

$$D_t \Theta = \frac{\Delta T}{H} \frac{\partial \Psi}{\partial x} + \kappa \nabla^2 \Theta.$$
(30)

Nondimensionalization of the problem yields equations including the dimensionless Prandtl number σ and the Rayleigh number R_a which are the control parameters of the problem. One can take the layer thickness H as the length of unit, the time $T = H^2/\kappa$ of vertical diffusion of heat as the unit of time, and the temperature difference ΔT as the unit of temperature.

Reduction of Dimensions and the Lorenz System

Saltzman (1962) derived a set of ordinary differential equations by expanding Ψ and Θ in double Fourier series in x and z, with functions of t alone for coefficients, and substituting these series into (29) and (30). A complete Galerkin approximation

$$\Psi(x,z,t) = \sum_{k=1}^{\infty} \sum_{l=1}^{\infty} \Psi_{k,l}(t) \sin\left(\frac{k\pi a}{H}x\right) \times \sin\left(\frac{l\pi}{H}z\right)$$
(31)

$$\Theta(x, z, t) = \sum_{k=1}^{\infty} \sum_{l=1}^{\infty} \Theta_{k,l}(t) \cos\left(\frac{k\pi a}{H}x\right) \times \sin\left(\frac{l\pi}{H}z\right)$$
(32)

yields an infinite set of ordinary differential equations for the time coefficients. He arranged the right-hand sides of the resulting equations in double Fourier series form, by replacing products of trigonometric functions of x (or z) by sums of trigonometric functions, and then equated coefficients of similar functions of x and z. He then reduced the resulting infinite system to a finite system by omitting reference to all but a specified finite set of functions of t. He then obtained time-dependent solutions by numerical integration. In certain cases, all, except three of the dependent variables, eventually tended to zero, and these three variables underwent irregular, apparently nonperiodic fluctuations. These same solutions would have been obtained if the series had been at the start truncated to include a total of three terms. Accordingly, in this study we shall let

$$\frac{a}{1+a^2}\kappa \quad \Psi = X\sqrt{2}\sin\left(\frac{\pi a}{H}x\right)\sin\left(\frac{\pi}{H}z\right)$$
(33)

$$\pi \frac{R_a}{R_c} \frac{1}{\Delta T} \quad \Theta = Y \sqrt{2} \cos\left(\frac{\pi a}{H}x\right) \sin\left(\frac{\pi}{H}z\right) - Z \sin\left(2\frac{\pi}{H}z\right)$$
(34)

where X(t), Y(t), and Z(t) are functions of time alone.

It is found that fields of motion of this form would develop if the Rayleigh number

$$R_a = \frac{g\alpha H^3 \Delta T}{\nu \kappa} \quad , \tag{35}$$

exceeds a critical value

$$R_c = \pi^4 a^{-2} (1+a^2)^3 \quad . \tag{36}$$

The minimum value of R_c , namely, $27\pi^4/4 = 657.51$, occurs when $a^2 = 1/2$. In fluid mechanics, the Rayleigh number for a fluid is a dimensionless number associated with the relation of buoyancy and viscosity in a flow. When the Rayleigh number is below the critical value for that fluid, heat transfer is primarily in the form of conduction; when it exceeds the critical value, heat transfer is primarily in the form of convection.

When the above truncation (33) and (34) is substituted into the dynamics, we obtain the equations (Lorenz model):

$$\dot{X} = -\sigma X + \sigma Y \tag{37}$$

$$\dot{Y} = rX - Y - XZ \tag{38}$$

$$\dot{Z} = -bZ + XY \tag{39}$$

Here a dot denotes a derivative with respect to the dimensionless time $t_d = \pi^2 H^{-2}(1 + a^2)\kappa t$, while $\sigma = \nu \kappa^{-1}$ is the Prandtl number, $r = R_a/R_c$, and $b = 4(1 + a^2)^{-1}$.

Fig. 6 Numerical solution of the Lorenz model, in the X - Y phase space with the parameters r = 28, $\sigma = 10$, and b = 8/3

Equations (37), (38), and (39) are called *Lorenz model* in the literature (Lorenz, 1960, 1963, 1984; Maas, 1994; Olbers, 2001). The system may give realistic results when the Rayleigh number is slightly supercritical, but their solutions cannot be expected to resemble those of the complete dynamics when strong convection occurs, in view of the extreme truncation. Figure 6 shows the numerical solution in the phase space with the parameters r = 28, $\sigma = 10$, and b = 8/3. The chaotic nature of this system inspired climate scientist and scientists in general. This phenomenon had probably the greatest impact of climate science to mathematics.

Scaling in the Climate System

As we will see now, the Coriolis effect is one of the dominating forces for the largescale dynamics of the oceans and the atmosphere. It is convenient to work in the rotating frame of reference of the Earth. The equation can be scaled by a length scale L, determined by the geometry of the flow, and by a characteristic velocity U. One can estimate the relative contributions in units of m/s^2 in the horizontal momentum equations:

$$\underbrace{\frac{\partial \mathbf{v}}{\partial t}}_{U/T \sim 10^{-8}} + \underbrace{\mathbf{v} \cdot \nabla \mathbf{v}}_{U^2/L \sim 10^{-8}} = \underbrace{-\frac{1}{\rho} \nabla p}_{\delta \mathbf{P}/(\rho \mathbf{L}) \sim 10^{-5}} + \underbrace{2\mathbf{\Omega} \times \mathbf{v}}_{\mathbf{f}_0 \mathbf{U} \sim 10^{-5}} + \underbrace{fric}_{\nu U/H^2 \sim 10^{-13}}$$
(40)

where fric denotes the contributions of friction due to eddy stress divergence (usually $\sim \nu \nabla^2 \mathbf{v}$). Typical values are given in Table 1. The values have been taken for the **ocean.**

It is furthermore useful to think about the orders of magnitude: Because of the continuity equation $U/L \sim W/H$ and since the horizontal scales are orders of magnitude larger than the vertical ones, the vertical velocity is very small relative to the horizontal. For small-scale motion (like small-scale ocean convection or cumulus clouds), the horizontal length scale is of the same order as the vertical one



	Quantity	Atmosphere	Ocean
Horizontal velocity	U	$10{\rm ms^{-1}}$	$10^{-1}{ m ms^{-1}}$
Vertical velocity	W	$10^{-1}{ m ms^{-1}}$	$10^{-4}{ m ms^{-1}}$
Horizontal length	L	10 ⁶ m	10 ⁶ m
Vertical length	Н	10 ⁴ m	10 ³ m
Horizonal pressure changes	δP (horizontal)	10 ³ Pa	10 ⁴ Pa
Mean pressure	P_0	10 ⁵ Pa	10 ⁷ Pa
Time scale	Т	10 ⁵ s	10 ⁷ s
Gravity (gravitation + centrifugal)	g	$10{\rm ms^{-2}}$	$10{\rm ms^{-2}}$
Earth radius	a	10 ⁷ m	10 ⁷ m
Coriolis parameter at 45°N	$f_0 = 2\Omega \sin \varphi_0$	$10^{-4} \mathrm{s}^{-1}$	$10^{-4} \mathrm{s}^{-1}$
2nd coriolis parameter at 45°N	$f_1 = 2\Omega \cos \varphi_0$	$10^{-4} \mathrm{s}^{-1}$	$10^{-4} \mathrm{s}^{-1}$
Density	ρ	$1 {\rm kg} {\rm m}^{-3}$	$10^3 \rm kg m^{-3}$
Viscosity (turbulent)	ν	$10^{-5} \mathrm{kg} \mathrm{m}^{-3}$	$10^{-6} \mathrm{kg} \mathrm{m}^{-3}$

Table 1 Table shows the typical scales in the atmosphere and ocean system. Using these orders of magnitude, one can derive estimates of the different terms in (40)

and therefore the vertical motion is in the same order of magnitude as the horizontal motion. The time scales are related to $T \sim L/U \sim H/W$.

It is essential to think about the relative importance of the different terms in the momentum balance (40). The Rossby Number Ro is the ratio of inertial (the left-hand side) to Coriolis (second term on the right-hand side) terms

$$Ro = \frac{(U^2/L)}{(fU)} = \frac{U}{fL}.$$
 (41)

It is used in the oceans and atmosphere, where it characterizes the importance of Coriolis accelerations arising from planetary rotation. It is also known as the Kibel number. Ro is small when the flow is in a so-called geostrophic balance.

Projection Methods: Coarse Graining and Stable Manifold Theory

The structure of fluid dynamical models and thus climate models is valid for systems with many degrees of freedom, many collisions, and for substances which can be described as a continuum. The transition from the highly complex dynamical equations to a reduced system is an important step since it gives more credibility to the approach and its results. The transition is also necessary since the active entangled processes are running on spatial scales from millimeters to thousands of kilometers, and temporal scales from seconds to millennia (Figs. 2 and 3). Therefore, the unresolved processes on subgrid scales have to be described. This is the typical problem in statistical physics, known as the so-called Mori-Zwanzig approach (Mori, 1965; Zwanzig, 1960, 1980). The basic idea is the evolution of

a system through a projection on a subset (macroscopic relevant part), where a randomness reflects the effects of the unresolved degrees of freedom. A particular example is the Brownian motion (Einstein, 1905; Langevin, 1908). Another solution for the transition form may degrees of freedom to the macroscopic laws goes back to Boltzmann (1896). The Boltzmann equation, also often known as the Boltzmann transport equation (Bhatnagar et al., 1954; Boltzmann, 1896; Cercignani, 1990), describes the statistical distribution of one particle in a fluid. It is one of the most important equations of nonequilibrium statistical mechanics, the area of statistical mechanics that deals with systems far from thermodynamic equilibrium. It is applied, for instance, when there is an applied temperature gradient or electric field. Both, the Mori-Zwanzig and Boltzmann approaches play also a fundamental role in physics. The microscopic equations show no preferred time direction, whereas the macroscopic phenomena in the thermodynamics have a time direction through the entropy. The underlying procedure is that part of the microscopic information is lost through coarse graining in space and time.

In order to get a first idea of coarse graining, one may think of the transition from Rayleigh-Bénard convection to the Lorenz system (section **Convection in the Rayleigh-Bénard system**). In our formula, the Galerkin approximation (31) and (32) provided a suitable projector to simply truncate the series at some specified wave number cutoff into a low-order system (such as in equations (33) and (34). The mathematical theory behind this truncation is called the center manifold theory (Haken, 1983; Oseledets, 1968). We could arrive at the slow manifold of the climate system, to which all the faster response variables (e.g., the atmosphere) are attracted. In mathematics, the slow manifold of an equilibrium point of a dynamical system occurs as the most common example of a center manifold. One of the main methods of simplifying dynamical systems is to reduce the dimension of the system to that of the slow manifold – center manifold theory rigorously justifies the modeling (Arnold, 1998; Arnold and Imkeller, 1998; Lorenz, 1986; Roberts, 2008).

The Mori-Zwanzig formalism (Mori, 1965; Zwanzig, 1960) and the slow manifold theory provide a conceptual framework for the study of dimension reduction and the parameterization of less relevant variables by a stochastic process. It includes a generalized (Langevin, 1908) theory. Langevin (1908) studied Brownian motion from a different perspective to Einstein's seminal 1905 paper (Einstein, 1905), describing the motion of a single Brownian particle as a dynamic process via a stochastic differential equation, as an Ornstein-Uhlenbeck process (Uhlenbeck and Ornstein, 1930).

The Gaussian filtering of hydrodynamic equations that leads to the Smagorinsky equations (Smagorinsky, 1963) is, in its essence, a version of coarse graining. The projection method includes the procedure to describe turbulent energy dissipation in turbulent flows, where the larger eddies extract energy from the mean flow and ultimately transfer some of it to the smaller eddies which, in turn, pass the energy to even smaller eddies, and so on up to the smallest scales, where the eddies convert the kinetic energy into internal energy of the fluid. At this scales (also known as Kolmogorov scale), the viscous friction dominates the flow (Frisch, 1996).

Brownian Motion, Weather, and Climate

The daily observed maximum and minimum temperatures are often compared to the "normal" temperatures based upon the 30-year average. Climate averages provide a context for something like "this winter will be wetter (or drier, or colder, or warmer, etc.) than normal." It has been said "Climate is what you expect. Weather is what you get." What is the difference between weather and climate? This can be also answered by an example/a metaphor in the football league. Predicting the outcome of the next game is difficult (weather), but predicting who will end up as German champion is unfortunately relatively easy (climate).

For climate, this transition between the climate and weather scales has been formulated conceptually (Hasselmann, 1976; Leith, 1975) and later re-formulated in a mathematical context (Arnold, 2001; Chorin et al., 1999; Gottwald, 2010). The effect of the weather on climate is seen by red-noise spectra in the climate system, showing one of the most fundamental aspects of climate and serving also as a null hypothesis for climate variability studies.

In a stochastic framework of climate theory, one may use an appropriate stochastic differential equation (Langevin equation)

$$\frac{d}{dt}x(t) = f(x) + g(x)\xi,$$
(42)

where $\xi = \frac{d}{dt}W(t)$ is a stationary stochastic process and the functions $f, g: \mathbb{R}^n \to \mathbb{R}^n$ describe the climate dynamics. The properties of the random force are described through its distribution and its correlation properties at different times. The process ξ is assumed to have a Gaussian distribution of zero average,

$$\langle \xi(t) \rangle = 0 \tag{43}$$

and to be δ -correlated in time,

$$\langle \xi(t)\xi(t+\tau) \rangle = \delta(\tau)$$
 (44)

where δ is the delta function defined by

$$\int_{\mathsf{R}} f(x)\,\delta(x-x_0)\,dx = f(x_0).$$
(45)

The brackets indicate an average over realizations of the random force. Formally, $\xi(t)$ is a random variable, i.e., $\xi(t)(\alpha)$ with different realizations due to random variable α . The expectation $\langle \xi(t) \rangle$ is thus the mean over all $\alpha :< \xi(t)(\alpha) \rangle_{\alpha}$. Using the ergodic hypothesis, the ensemble average $\langle \rangle$ can be expressed as the time average $\lim_{T\to\infty} \frac{1}{T} \int_{-T/2}^{T/2} dt$ of the function. Almost all points in any subset of the phase space eventually revisit the set. For a Gaussian process, only the average and second moment need to be specified since all higher moments can be expressed in terms of the first two. Note that the dependence of the correlation function on the

time difference τ assumes that ξ is a stationary process. ξ is called a white-noise process.

Additionally, there might be an external forcing F(x, t) which is generally time, variable, and space dependent. In his theoretical approach, Hasselmann (1976) formulated a linear stochastic climate model

$$\frac{d}{dt}x(t) = Ax + \sigma\xi + F(t), \tag{46}$$

with system matrix $A \in \mathbb{R}^{n \times n}$, constant noise term σ , and stochastic process ξ . Many features of the climate system can be well described by (46), which is analogous to the Ornstein-Uhlenbeck process in statistical physics (Uhlenbeck and Ornstein, 1930). Notice that $\sigma\xi$ represents a stationary random process. The relationship derived above is identical to that describing the diffusion of a fluid particle in a turbulent fluid. In a time scale separated system, during one slow-time unit the fast uninteresting variables y perform many "uncorrelated" events (provided that the fast dynamics are sufficiently chaotic). The contribution of the uncorrelated events to the dynamics of the slow interesting variables x is as a sum of independent random variables. By the weak central limit theorem, this can be expressed by a normally distributed variable. Note, in the absence of any feedback effects Ax, the climate variations would continue to grow indefinitely as the Wiener process. A perturbation in a system with a negative feedback mechanism will be reduced, whereas in a system with positive feedback mechanisms, the perturbation will grow. In the one-dimensional case, A can be rewritten as $-\lambda$. The real part of λ determines then the stability of the system and is called feedback factor.

Climate Variability and Sensitivity

Imagine now that the temperature of the ocean mixed layer of depth h is governed by a one-dimensional system

$$\frac{dT}{dt} = -\lambda T + Q_{\text{net}} + f(t), \qquad (47)$$

where the air-sea fluxes due to weather systems are represented by a white-noise process with zero average $\langle Q_{\text{net}} \rangle = 0$ and δ -correlated in time $\langle Q_{\text{net}}(t)Q_{\text{net}}(t + \tau) \rangle = \delta(\tau)$. The function f(t) is a time-dependent deterministic forcing. Assume furthermore that $f(t) = c \cdot u(t)$ with u(t) as unit step or the so-called Heaviside step function. Because $\langle Q_{\text{net}} \rangle = 0$, $\langle T(t) \rangle$ can be solved using the Laplace transform:

$$\langle T(t) \rangle = \mathcal{L}^{-1}\{F(s)\}(t) = \mathcal{L}^{-1}\left\{\frac{\langle T(0) \rangle}{s+\lambda} + \frac{c}{s} \cdot \frac{1}{s+\lambda}\right\}$$
(48)

$$= T(0) \cdot \exp(-\lambda t) + \frac{c}{\lambda} (1 - \exp(-\lambda t))$$
(49)

because we have $\langle T(0) \rangle = T(0)$. As equilibrium response, we have

$$\Delta T = \lim_{t \to \infty} \langle T(t) \rangle = \frac{c}{\lambda} .$$
(50)

The fluctuation can be characterized by the spectrum

$$S(\omega) = \langle \hat{T}\hat{T}^* \rangle = \frac{1}{\lambda^2 + \omega^2}.$$
(51)

and therefore, the spectrum and the equilibrium response are closely coupled (fluctuation-dissipation theorem). In mathematics, this is called Wiener-Chintschin-Theorem (Chintchin, 1934; Wiener, 1930). For some energy considerations, it is useful to rewrite equation (47) as

$$C \frac{dT}{dt} = -\lambda_C T + f_C , \qquad (52)$$

with $C = c_p \rho dz$ as the heat capacity of the ocean. For a depth of 200 m of water distributed over the globe, $C = 4.2 \cdot 10^3 \text{Ws} \text{kg}^{-1} \text{K}^{-1} \times 1000 \text{kg} \text{m}^{-3} \times 200 \text{m} = 8.4 \cdot 10^8 \text{Ws} \text{m}^{-2} \text{K}^{-1}$. The temperature evolution is

$$T(t) = T(0) \cdot \exp(-\lambda_c/C t) + \frac{f_C}{\lambda_C} \left(1 - \exp(-\lambda_C/C t)\right)$$
(53)

The left-hand side of (52) represents the heat uptake by the ocean, which plays a central role in the transient response of the system to a perturbation (53).

Typical changes in f_C are 4 W m⁻² for doubling of CO_2 , $\lambda_C = 1 - 2 \text{ Wm}^{-2}\text{K}^{-1}$. The typical time scale for a mixed layer ocean is $C/\lambda_C = 13 - 26$ years. Please note that the climate system is simplified by a slab ocean with homogeneous temperature and heat capacity. This is an approximation as the heat capacity should vary in time as the perturbation penetrates to deeper oceanic levels. The equilibrium temperature change ΔT is

$$\Delta T = \frac{\Delta f_C}{\lambda_C} = \frac{c}{\lambda} \tag{54}$$

with values of $\Delta T = 2 - 4$ K. The term $CS = \frac{1}{\lambda_C}$ is called climate sensitivity to a radiative forcing Δf_C :

$$\Delta T = CS \cdot \Delta f_C. \tag{55}$$

In the literature, the concept of climate sensitivity is quite often used as the equilibrium temperature increase for a forcing Δf_C related to doubling of CO_2 . It is obvious that the CS depends on the included sources of feedback of the system which are related to climate components and their respective time scales (e.g., Lohmann 2018). Due to the non-normality in (46), the effective damping may not be directly related to the eigenvalues of the system.

Non-normal Growth of the Climate System

In the one-dimensional case for $x(t) = \exp(at)$, we have the inverse Laplace transform

$$\exp(at) = \mathcal{L}^{-1}\{F(s)\}(t) = \frac{1}{2\pi i} \lim_{T \to \infty} \int_{\gamma - iT}^{\gamma + iT} e^{st} \frac{1}{s - a} \, ds, \tag{56}$$

and the entire range of t is controlled t by the resolvent $\left|\frac{1}{s-a}\right|$. Using the Fourier transformation, (46) with forcing F(t) is transformed to

$$(i\omega I - A)\hat{x} = \hat{F} \tag{57}$$

$$\hat{x} = (i\omega I - A)^{-1}\hat{F} \tag{58}$$

where I is the identity. The so-called resolvent operator of matrix A is $R(\omega) = (i\omega I - A)^{-1}$. The behavior of the norms $||\exp(At)||$ over the entire range of t is controlled t by the resolvent norm $||R(\omega)||$. If A is a normal operator

$$A A^+ = A^+ A, (59)$$

where ⁺ denotes the adjoint-complex operator, then

$$||R(\omega)|| = 1/\operatorname{dist}(i\omega, \sigma(A)) \tag{60}$$

is completely determined by the spectrum $\sigma(A)$ alone. The operator dist denotes the shortest distance of ω to the eigenvalues, the spectrum $\sigma(A)$. This explains the success of eigenvalue analysis. In contrast to this, for non-normal operators the behavior of $||R(\omega)||$ may deviate from that dramatically, and hence in this context pseudospectral analysis is just the right tool. For example, there are problems in fluid mechanics where $\sigma(A)$ is contained in the left half-plane, which suggests laminar behavior, but it protrudes strongly into the right half-plane, which implies that $||e^{At}||$ has a big hump before decaying exponentially fast to zero Reddy et al. (1993) and Trefethen et al. (1993). More about the dynamics can be learned by examining the pseudospectrum of A in the complex plane. Inspection of many geophysical systems shows that most of the systems fail the normality condition. The ϵ -pseudospectrum of operator A is defined by two equivalent formulations:

$$\Lambda_{\epsilon}(A) = \{ z \in \mathbb{C} : ||(zI - A)^{-1}|| \ge \epsilon^{-1} \}$$

= $\{ z \in \mathbb{C} : [$ smallest singular value of $(zI - A)] \le \epsilon \}.$ (61)

This set of values z in the complex plane are defined by contourlines of the resolvent $(zI - A)^{-1}$. The resolvent determines the system's response to a forcing as supplied by external forcing F(x, t), stochastic forcing $g(x)\xi$, or initial/boundary conditions. The pseudospectrum reflects the robustness of the spectrum and provides information about instability and resonance. One theorem is derived from Laplace transformation stating that transient growth is related to how far the ϵ -pseudospectrum extends into the right half-plane:

$$||\exp(At)|| \ge \frac{1}{\epsilon} \sup_{z \in A_{\epsilon}(A)} \operatorname{Real}(z).$$
 (62)

In terms of climate theory, the pseudospectrum indicates resonant amplification. Maximal amplification is at the poles of $(zI - A)^{-1}$, characterized by the eigenfrequencies. In a mathematical normal matrix A, the system's response is characterized solely by the proximity to the eigenfrequencies. In the non-normal case, the pseudospectrum shows large resonant amplification for frequencies which are not eigenfrequencies. This transient growth mechanism is important for both initial value and forced problems.

An atmospheric general circulation model PUMA (Fraedrich et al., 2005) is applied to the problem. The model is based on the multilevel spectral model described by Hoskins and Simmons (1975). For our experiments we chose five vertical levels and a T21 horizontal resolution. PUMA belongs to the class of models of intermediate complexity (Claussen et al., 2002); it has been used to understand principle feedbacks (Lunkeit et al., 1998), and dynamics on long time scales (Romanova et al., 2006). For simplicity, the equations are scaled here such that they are dimensionless. The model is linearized about a zonally symmetric mean state providing for a realistic storm track at midlatitudes (Frisius et al., 1998). In a simplified version of the model and calculating the linear model A with n = 214, one can derive the pseudospectrum. Figure 7 indicates resonances besides the poles (the eigenvalues) indicated by crosses. The Im(z)-axis shows the frequencies and the Re(z)-axis the damping/amplification of the modes. Important modes for the climate system are those with -0.5 < Im(z) < 0.5 representing planetary Rossby waves. The basic feature is that transient growth of initially small perturbations can occur even if all the eigenmodes decay exponentially. Mathematically, an arbitrary matrix A can be decomposed as a sum

$$A = D + N \tag{63}$$

where A is diagonalizable and N is nilpotent (there exists an integer $q \in N$ with $N^q = 0$) and D commutes with N (i.e., DN = NA). This fact follows from the Jordan-Chevalley decomposition theorem. This means that we can compute the



Fig. 7 Contours of $\log_{10}(1/\epsilon)$. The figure displays resonant structures of the linearized atmospheric circulation model. The modes extend to the right half-plane and are connected through resonant structures, indicating for transient growth mechanism inherent in atmospheric dynamics

exponential of (A t) by reducing to the cases:

$$\exp(At) = \exp((D+N)t) = \exp(Dt) \quad \exp(Nt) \tag{64}$$

where the exponential of Nt can be computed directly from the series expansion, as the series terminates after a finite number of terms. Basically, the number $q \in N$ is related to the transient growth of the system (q = 1 means no transient growth).

The resonant structures are due to the mode interaction: It is not possible to change one variable without the others, because they are not orthogonal. Interestingly, one can also compute the A^+ model, showing the optimal perturbation of a mode through its biorthogonal vector which is the associated eigenvector of the adjoint A^+ . The analysis indicates that non-normality of the system is a fundamental feature of the atmospheric dynamics. This has consequences for the error growth dynamics and instability of the system, e.g. Palmer (1996) and Lohmann and Schneider (1999). Similar features are obtained in shear flow systems (Reddy et al., 1993; Trefethen et al., 1993) and other hydrodynamic applications. This transient growth mechanism is important for both initial value and forced problems of the climate system (Farrell and Ioannou, 1996).

Predictability

In climate we may ask about our initial state. Climatologists always feel uncertain when we want to give the initial values. There is always a more or less big inaccuracy due to weather and uncertainties in many quantities which cannot be observed at any time step (e.g., the deep ocean). The Lyapunov exponents λ play an important role in knowing the predictability of a system. This is because the larger they are, the smaller the number of steps for which predictions can be made with a certain, desired accuracy. Consider a trajectory x(t) and a nearby trajectory $x(t) + \delta(t)$ where $\delta(t)$ is a vector with infinitesimal initial length. As the system evolves, track how $\delta(t)$ changes. The maximal Lyapunov exponent of the system is the number λ such that $|\delta(t)| \approx |\delta(0)| \cdot \exp(\lambda t)$. A classical example is again the Lorenz system (37), (38), and (39) where for large parts of the phase space, we have limited predictability because initial errors can grow.

Every dynamical system has a spectrum of Lyapunov exponents, one for each dimension of its phase space. Like the largest eigenvalue of a matrix, the largest Lyapunov exponent is responsible for the dominant behavior of a system. In case of weather and climate, this Lyapunov exponent is therefore also time scale dependent. Causality in climate has only a limited range in time and can only be verified in the context of finite errors. Please note that even in classical mechanics, strict causal relationships cannot be verified experimentally! It is true that the classical laws of motion are generally deterministic. But the connection with the real physical world is always possible only with limited accuracy due to the unavoidable measurement errors. Therefore, the actual state can only be given with a certain probability distribution within state ranges. In the usual discussion, this important part of physics is often faded out; one likes to limit oneself to the equations of motion alone.

Natural events also have their own time scales, namely, the so-called Lyapunov times t_{Lyap} (these result from the expansion rates at $t_{Lyap} = \lambda^{-1}$). This Lyapunov time of a weather situation, it is about seconds to days, for climate years to decades. We now have to compare the two relevant time scales t_M and t_{Lyap} . Three cases are possible:

- The Lyapunov time of the climate system under investigation is much larger than the humanly relevant time scale ($t_{Lyap} >> t_M$): Then we can imagine the initial state inaccuracy to be smaller and smaller in our minds, because this would only increase the prediction time. Even if it would become infinite in our thoughts because we let the measurement inaccuracy become zero, we would not even notice it. Therefore, in these cases we consider the event (within the accepted accuracy) to be predictable, causal.
- The human relevant time scale is much larger than the Lyapunov time of the investigated system ($t_M >> t_{Lyap}$): In this case prediction is no longer possible; the actual course is completely different from the expected one. We observe statistical, random behavior, since we do not know the actual initial state.

• The relevant time scale is about as large as the Lyapunov time of the system under study ($t_M \approx t_{Lyap}$): Then no exact but approximate predictions are possible; it is also not entirely random, statistically. The predictions can even be improved by measurement progress or by less demanding requirements for prediction accuracy. A good example is the weather forecasts, which are only possible to a limited extent; in the short and medium term, they are now quite reliable.

In essence, therefore, causality depends on the time of interest in comparison to the forecast time, whether we can regard a phenomenon "practically" as causal, as predictable in (sometimes excellent approximation), or whether the event appears to us to be completely random, or finally as lying in the transition area and is therefore experienced as improvable by increasing the accuracy, i.e., as neither causal nor statistical. Climate is not only a differential equation; it must also be coupled to the real world by specifying the initial values with measurement errors and by translating the final values into measurable predictions. Thus it loses its purely mathematical, causal character determined by the solution of differential equations.

Besides the initial conditions, uncertainties can appear through the external forcing. Prominent external forcing are the change in greenhouse gases into the atmosphere (e.g., Fig. 8) which strongly affect the long-term evolution of the Earth system. Another external forcing is due to changes in insolation by orbital parameters. There parameters vary on multi-millennial time scales (thousand years = ky) and can be calculated by orbital theory. Milankovitch (1941) suggested the ice sheet growth and decay is triggered by this external forcing. There are several open questions for paleoclimate dynamics. Despite the pronounced change in Earth system response evidenced in paleoclimatic records, the frequency and amplitude characteristics of the orbital parameters, i.e., eccentricity ($\sim 100 \text{ ky}$), obliquity $(\sim 41 \text{ ky})$, and precession $(\sim 21 \text{ and } \sim 19 \text{ ky})$, do not vary (Berger and Loutre, 1991), the climate frequency does. The uncertainty on long time scale is usually dominated by the external forcing and the short time scale by the initial value problem; the intermediate times at 10–50 years for the coming decades are dominated by internal variability and uncertainty in model physics (Hawkins and Sutton, 2009). The uncertainty of global and especially regional temperature estimates on decadal to multidecadal time scales is manifested by large-scale coherent pattern like AMO, PDO, and the quasi-decadal mode (cf. Fig. 2). For a while, people tended to think that deterministic models would still always provide the best models in these cases. One thought of the climate system is that deterministic models would be completely adequate for describing the Earth's atmosphere, which is basically just a layer of gas subject to external heating. As seen here, the phenomenon of stochasticity means that this is not so. As in the Lorenz system, chaos is introduced into the climate system. So, in many cases, for quite fundamental reasons, deterministic mathematical models do not provide adequate models. Statistical models are mathematical models, each replicate realization of which will be different from other realizations with the same model, even under identical conditions. Statistical models are the major means of making sense of the climate dynamics (Fig. 9).



Fig. 8 Different climate scenarios with a coupled Earth system model (Ackermann et al., 2020). Time series of 11-year mean (**a**) CO_2 forcing as concentration of CO_2 equivalent in the atmosphere, (**b**) global near-surface average temperature, (**c**) sea ice volume in the Northern Hemisphere; shaded areas indicate 1 standard deviation

Boltzmann Dynamics

One of the most significant theoretical breakthroughs in statistical physics was due to Ludwig Boltzmann (Boltzmann 1896, 1995 for a recent reprint of his famous lectures on kinetic theory), who pioneered nonequilibrium statistical mechanics. Boltzmann postulated that a gas was composed of a set of interacting particles, whose dynamics could be (at least in principle) modeled by classical dynamics. Due to the very large number of particles in such a system, a statistical approach was adopted, based on simplified physics composed of particle streaming in space and billiard-like inter-particle collisions (which are assumed elastic).

As already mentioned above, a fluid can be described by several physical theories, of different granularities. The fact that we can, in principle, recover the



phenomena predicted by the coarse-grained theories from solutions of the finegrained theories also suggests a non-conventional way of constructing numerical algorithms for simulating fluid flows: Instead of directly modeling the coarsegrained equations (i.e., Navier-Stokes equations for human-scale flows), we can construct a simplified model of the fine-grained equations, which will exhibit the same behavior at the larger scales.

In the following, an example derived for the Lattice Boltzmann Model (LBM) is shown which is related to the thermohaline circulation. Water that is dense enough to sink from the surface to the bottom is formed when cold air blows across the ocean at high latitudes in winter in the northern North Atlantic (e.g., in the Labrador Sea and between Norway and Greenland) and near Antarctica. The wind cools and evaporates water. If the wind is cold enough, sea ice forms, further increasing the salinity of the water because sea ice is fresher than seawater and salty water remains in the water when ice is formed. Bottom water is produced only in these regions, and the deep ocean is affected by these deepwater formation processes. In other regions, cold, dense water is formed, but it is not quite salty enough to sink to the bottom. At mid and low latitudes, the density, even in winter, is sufficiently low that the water cannot sink more than a few hundred meters into the ocean. The only exception are some seas, such as the Mediterranean Sea, where evaporation is so great that the salinity of the water is sufficiently great for the water to sink to intermediate depths in the seas. If these seas are can exchange water with the open ocean, the waters formed in winter in the seas spreads out to intermediate depths in the ocean. A numerical solution of this equation is shown in Fig. 10.





Conclusions

Climate change occurred during the history of the Earth, the tectonic movements over billions of years, and climate has varied between extremes before any anthropogenic action could have arisen. However, anthropogenic action in terms of heavy usage of fossil fuel has the potential to affect the Earth to a point where its habitability is significantly affected. In terms of the time scale, it is noted that we might disturb planetary-scale processes in the course of a few decades. The complication is due to the fact that the climate system has inherent fluctuations (internal climate variability), uncertainties in model formulations, and scenario uncertainties for past and future climate scenarios. Modeling is necessary to produce a useful understanding of abrupt climate processes. Model analyses help to focus research on possible causes of abrupt climate change, such as human activities; on key areas where climatic thresholds might be crossed; and on fundamental uncertainties in climate system dynamics. Improved understanding of abrupt climatic changes that occurred in the past and that are possible in the future can be gained through climate models. For climate science, most fundamental laws were discovered decades ago (Landau and Lifshitz, 1987) (although there are Navier-Stokes existence and smoothness problems in three dimensions, see The Clay Mathematics Institute, http://www.claymath.org/millennium-problems/). The system has specific scales and characteristic numbers. Part of the uncertainty is due to the difficulty to find a proper description of the system. Since it is hard to disagree with the simple statement "more data are better," the task here is rather to identify those dimensions in the data space where invested resources may yield to a maximum of new information. In this way, data assimilation techniques could help for an estimate of the state of the system but also its uncertainty (Burgers et al., 1998; Kalman, 1960; Nerger and Hiller, 2013).

High-resolution models are required to elucidate the causal chains in the climate system, notably during abrupt transitions of the last deglaciation, and provide a benchmark for future transitions under rapid CO_2 increase. Practically, given the present high-performance computer capacities, efficient and parallelized model codes, it is now possible to conduct simulations for 50–100 model years per day even with a multi-scale ansatz (Lohmann et al., 2020; Sein et al., 2018). Recent developments have considerably improved the computational efficiency and scalability of unstructured-mesh approaches on high-performance computing systems (Danilov et al., 2017). The surface ocean current in such high-resolution simulation (Fig. 11) has a completely different structure including eddies than the structure in coarse resolution model.

Weather and climate extremes cause huge economic damages and harm many lives each year (Franzke, 2017). There is evidence that some types of weather and climate extremes, like heat waves and flooding, have already increased or intensified over the last few decades, and climate projections reveal a further intensification for many types of weather and climate extremes in many regions though the uncertainties still remain large. Future research may be enhanced along

Fig. 11 Representation of small-scale features of ocean currents in high-resolution ocean models: simulated velocity field: simulated velocity field in the North Atlantic at 100 m depth in December 1950 using FESOM with high-resolution locally eddy-resolving mesh based on Sein et al. (2018)



three directions data, statistics, theory, and models, leading to an increase in the current knowledge about the climate evolution. It is crucial that researchers deepen or acquire the ability to integrate all directions into their arsenal of mathematical methods.

Cross-References

- Interdisciplinary Mathematics and Sciences in Schematic Ocean Current Maps in the Seas Around Korea
- Limit Cycles in Planar Systems of Ordinary Differential Equations

References

- Ackermann L, Danek C, Gierz P, Lohmann G (2020) Amoc recovery in a multicentennial scenario using a coupled atmosphere-ocean-ice sheet model. Geophys Res Lett 47(16):e2019GL086,810 Arnold L (1998) Random dynamical systems. Springer, Berlin/Heidelberg
- Arnold L (2001) Hasselmann's program revisited: the analysis of stochasticity in deterministic climate models, vol 49. Birkhäuser, Boston
- Arnold L, Imkeller P (1998) Normal forms for stochastic differential equations. Probab Theory Related Fields 110(4):559–588
- Berger A, Loutre MF (1991) Insolation values for the climate of the last 10 million years. Quat Sci Rev 10(4):297–317
- Bhatnagar P, Gross EP, Krook MK (1954) A model for collision process in gases. I. Small amplitude processes in charged and neutral one-component system. Phys Rev 94:511

- Boltzmann L (1896) Vorlesungen über Gastheorie: 2 volumes (in German). Leipzig 1895/98 UB: O 5262–6.
- Boltzmann L (1995) Lectures on gas theory. Dover Publications, New York. ISBN:978-0486684550
- Buckingham E (1914) On physically similar systems; illustrations of the use of dimensional equations. Phys Rev 4(4):345–376. https://doi.org/10.1103/PhysRev.4.345
- Burgers G, Jan van Leeuwen P, Evensen G (1998) Analysis scheme in the ensemble kalman filter. Mon Weather Rev 126(6):1719–1724
- Cercignani C (1990) Mathematical methods in kinetic theory, 2nd edn. Plenum. ISBN:978-0306434600
- Chintchin A (1934) Korrelationstheorie der stationären stochastischen prozesse. Math Ann 109(1):604-615
- Chorin AJ, Kast AP, Kupferman R (1999) Unresolved computation and optimal predictions. Commun Pure Appl Math 52(10):1231–1254
- Claussen M, Mysak L, Weaver A, Crucifix M, Fichefet T, Loutre MF, Weber S, Alcamo J, Alexeev V, Berger A, et al (2002) Earth system models of intermediate complexity: closing the gap in the spectrum of climate system models. Clim Dyn 18(7):579–586
- Danilov S, Sidorenko D, Wang Q, Jung T (2017) The finite-volume sea ice–ocean model (fesom2). Geosci Model Dev 10:765–789
- Einstein A (1905) Investigations on the theory of the brownian movement. Ann der Phys 17:549– 560
- Farrell BF, Ioannou PJ (1996) Generalized stability theory. Part I: autonomous operators. J Atmos Sci 53(14):2025–2040
- Fraedrich K, Jansen H, Kirk E, Luksch U, Lunkeit F (2005) The planet simulator: towards a user friendly model. Meteorol Z 14(3):299–304
- Franzke CL (2017) Impacts of a changing climate on economic damages and insurance. Econ Disasters Clim Change 1(1):95–110
- Frisch U (1996) Turbulence: the legacy of A.N. Kolmogorov. Cambridge University Press. ISBN:0-521-45103-5
- Frisius T, Lunkeit F, Fraedrich K, James IN (1998) Storm-track organization and variability in a simplified atmospheric global circulation model. Q J R Meteorol Soc 124(548):1019–1043
- Gottwald G (2010) On recent trends in climate dynamics. AMS Gazette 37(5)
- Haken H (1983) Synergetics. Springer, Berlin/Heidelberg
- Hasselmann K (1976) Stochastic climate models. Part I. Theory. Tellus 6:473-485
- Hawkins E, Sutton R (2009) The potential to narrow uncertainty in regional climate predictions. Bull Am Meteorol Soc 90(8):1095–1108
- Hoskins B, Simmons A (1975) A multi-layer spectral model and the semi-implicit method. Q J R Meteorol Soc 101(429):637–655
- Kalman RE (1960) A new approach to linear filtering and prediction problems. J Basic Eng 82(1):35–45. https://doi.org/10.1115/1.3662552
- Landau LD, Lifshitz EM (1959) Fluid mechanics, course of theoretical physics, vol 6. Pergamon Press, Oxford
- Landau LD, Lifshitz EM (1987) Course of theoretical physics: fluid mechanics, vol 6, 2nd edn. Butterworth-Heinemann. ISBN:978-0750627672
- Langevin P (1908) On the theory of brownian motion. Comptes Rendues 146:530-533
- Leith C (1975) Climate response and fluctuation dissipation. J Atmos Sci 32(10):2022–2026
- Lisiecki LE, Raymo ME (2005) A pliocene-pleistocene stack of 57 globally distributed benthic 180 records. Paleoceanography 20(1). https://doi.org/10.1029/2004pa001071
- Lohmann G (2018) ESD ideas: the stochastic climate model shows that underestimated holocene trends and variability represent two sides of the same coin. Earth Syst Dyn 9(4):1279–1281
- Lohmann G, Schneider J (1999) Dynamics and predictability of stommel's box model. A phasespace perspective with implications for decadal climate variability. Tellus A 51(2):326–336
- Lohmann G, Butzin M, Eissner N, Shi X, Stepanek C (2020) Abrupt climate and weather changes across time scales. Paleoceanogr Paleoclimatol 35(9). https://doi.org/10.1029/2019pa003782

Lorenz EN (1960) Maximum simplification of the dynamic equations. Tellus 12(3):243-254

- Lorenz EN (1963) Deterministic nonperiodic flow. J Atmos Sci 20(2):130-141
- Lorenz EN (1984) Irregularity: a fundamental property of the atmosphere*. Tellus A 36(2):98–110 Lorenz EN (1986) On the existence of a slow manifold. J Atmos Sci 43(15):1547–1558
- Lunkeit F, Fraedrich K, Bauer S (1998) Storm tracks in a warmer climate: sensitivity studies with a simplified global circulation model. Clim Dyn 14(11):813–826
- Maas LR (1994) A simple model for the three-dimensional, thermally and wind-driven ocean circulation. Tellus A 46(5):671–680
- Milankovitch MK (1941) Kanon der erdbestrahlung und seine anwendung auf das eiszeitenproblem. R Serb Acad Spec Publ 133:1–633
- Mori H (1965) Transport, collective motion, and brownian motion. Prog Theor Phys 33(3):423-455
- Morice CP, Kennedy JJ, Rayner NA, Jones PD (2012) Quantifying uncertainties in global and regional temperature change using an ensemble of observational estimates: the hadCRUT4 data set. J Geophys Res Atmos 117(D8)
- Nerger L, Hiller W (2013) Software for ensemble-based data assimilation systems–implementation strategies and scalability. Comput Geosci 55:110–118
- Olbers D (2001) A gallery of simple models from climate physics. In: Imkeller P, von Storch J (eds) Stochastic climate models, progress in probability, 49:3–63
- Oseledets VI (1968) A multiplicative ergodic theorem. Characteristic Ljapunov, exponents of dynamical systems. Trudy Moskovskogo Matematicheskogo Obshchestva 19:179–210
- Palmer T (1996) Predictability of the atmosphere and oceans: from days to decades. In: Decadal climate variability. Springer, Berlin/Heidelberg, pp 83–155
- Peixoto JP, Oort AH (1992) Physics of climate. American Institute of Physics, New York
- Rayleigh L (1916) On convection currents in a horizontal layer of fluid, when the higher temperature is on the under side. Phil Mag 6:529–546
- Reddy SC, Schmid PJ, Henningson DS (1993) Pseudospectra of the orr-sommerfeld operator. SIAM J Appl Math 53(1):15–47
- Roberts AJ (2008) Normal form transforms separate slow and fast modes in stochastic dynamical systems. Phys A Stat Mech Appl 387(1):12–38
- Romanova V, Lohmann G, Grosfeld K, Butzin M (2006) The relative role of oceanic heat transport and orography on glacial climate. Quat Sci Rev 25(7–8):832–845
- Saltzman B (1962) Finite amplitude free convection as an initial value problem I. J Atmos Sci 19:329–341
- Sein DV, Koldunov NV, Danilov S, Sidorenko D, Wekerle C, Cabos W, Rackow T, Scholz P, Semmler T, Wang Q, et al (2018) The relative influence of atmospheric and oceanic model resolution on the circulation of the north atlantic ocean in a coupled climate model. J Adv Model Earth Syst 10(8):2026–2041
- Smagorinsky J (1963) General circulation experiments with the primitive equations: I. The basic experiment. Mon Weather Rev 91(3):99–164
- Trefethen LN, Trefethen AE, Reddy SC, Driscoll TA (1993) Hydrodynamic stability without eigenvalues. Science 261(5121):578–584
- Uhlenbeck GE, Ornstein LS (1930) On the theory of the brownian motion. Phys Rev 36(5):823
- Wiener N (1930) Generalized harmonic analysis. Acta Math 55:117-258
- Zwanzig R (1960) Ensemble method in the theory of irreversibility. J Chem Phys 33:1338
- Zwanzig R (1980) Problems in nonlinear transport theory. In: Systems far from equilibrium. Springer, Berlin/Heidelberg, pp 198–225