

Climate System II

(Winter 2023/2024)

6th lecture:

Biogeochemical cycles, vegetation and dust

(Aridity and dust, vegetation dynamics, land use, terrestrial biosphere)


Gerrit Lohmann, Martin Werner

Tuesday, 10:15-11:45


https://paleodyn.uni-bremen.de/study/climate2023_24.html

What are biogeochemical cycles?

- **Earth system has four parts**
 - *atmosphere*
 - *hydrosphere*
 - *lithosphere*
 - *biosphere*
- **Biogeochemical cycles**
 - *The chemical interactions (cycles) that exist between the atmosphere, hydrosphere, lithosphere, and biosphere*
- **Abiotic (physio-chemical) and biotic processes drive these cycles**




Transport of biogeochemical aerosols and dust



Animation 1. Aerosol optical thickness of black and organic carbon (green), dust (red-orange), sulfates (white), and sea salt (blue) from a 10 km resolution GEOS-5 "nature run" using the GOCART model. The animation shows the emission and transport of key tropospheric aerosols from August 17, 2006 to April 10, 2007.

Biogeochemical cycles in climate research



**Biogeochemical cycles are a key element
for understanding our past and present climate!**

Biogeochemical cycles - key elements

Six nutrient elements make up 95% of the biomass mass on earth and form the biochemical foundation for life.

- **Carbon (CO_2 , CH_4 , CO)**
- **Nitrogen (N_2O , NO , NO_2 , NH_3)**
- **Sulfur (SO_2 , COS , H_2S , H_2SO_4)**
- **Phosphorous**
- **Hydrogen**
- **Oxygen**
- **Water**

Biogeochemical cycles: Common features of all key elements


- **each element typically occurs in all four parts of the Earth System (e.g. water, carbon, nitrogen, etc.)**
- **each biogeochemical cycle can be described by**
 - ***reservoirs A***
 - ***fluxes F in and out of pools***
 - ***chemical or biochemical transformations***
 - ***important quantity: turnover times τ***
 - ***τ can be calculated as the size of reservoir A divided by the sum of all ingoing (or outgoing) fluxes F***

Biogeochemical cycles - key elements

Six nutrient elements make up 95% of the biomass mass on earth and form the biochemical foundation for life.


- **Carbon (CO₂, CH₄, CO)**
- **Nitrogen (N₂O, NO, NO₂, NH₃)**
- **Sulfur (SO₂, COS, H₂S, H₂SO₄)**
- **Phosphorous**
- **Hydrogen**
- **Oxygen**
- **Water**

The global carbon cycle



⇒ PEP-lecture: Carbon Cycle

CO₂ sources and sinks



© Global Carbon Project • Data: CDIAC/GCP/NOAA-ESRL/UNFCCC/BP/USGS

[[<https://www.globalcarbonproject.org/carbonbudget/19/presentation.htm>]]

(percentage for years 2009-2018)

Carbon isotopes

■ Relative abundance


- ^{12}C 98.89 % **stable**
- ^{13}C 1.11 % **stable**
- ^{14}C 1×10^{-10} % **half-life=5276 years**

■ ^{12}C and ^{13}C are stable isotopes

■ ^{14}C is called radiocarbon

- Formed by cosmic radiation ($^{14}\text{N} + ^1_0\text{n} \rightarrow ^{14}\text{C} + ^1_1\text{p}$)
 - Also formed in nuclear explosions
-
- *During photosynthesis, plants preferentially uptake ^{12}C
=> the atmosphere will become enriched in ^{13}C*


Variations of ^{13}C in different carbon pools




Variations of ^{13}C due to fossil fuel burning

Mauna Loa Observatory, Hawaii and South Pole, Antarctica Monthly Average Carbon Isotopic Trends

Data from Scripps CO₂ Program Last updated March 2009




Variations of ^{13}C since the last glacial maximum (LGM)




[Brook, The Ice Age Carbon Puzzle, Science, 2012]

The global nitrogen cycle



The global nitrogen cycle. Pools (\square) and annual (\rightarrow) flux in 10^{12} gN_2 . Note that the industrial fixation of nitrogen is nearly equal to the global biological fixation. (SOURCE: Data from Söderlund, and T. Rosswall, 1982, O. Hutzinger (ed.), *The Handbook of Environmental Chemistry*, Vol 1, Pt. B., Springer-Verlag New York, Inc., New York).

The global sulfur cycle




Biogeochemical cycles - key elements

Six nutrient elements make up 95% of the biomass mass on earth and form the biochemical foundation for life.

- **Carbon (CO_2 , CH_4 , CO)**
- **Nitrogen (N_2O , NO , NO_2 , NH_3)**
- **Sulfur (SO_2 , COS , H_2S , H_2SO_4)**
- **Phosphorous**
- **Hydrogen**
- **Oxygen**
- **Water**

Dust in the climate system



Dust depositions on glacial-interglacial time scales

Dust–climate couplings over the past 800,000 years from the EPICA Dome C ice core

Vol 452 | 3 April 2008 | doi:10.1038/nature06763

F. Lambert^{1,2}, B. Delmonte³, J. R. Petit⁴, M. Bigler^{1,5}, P. R. Kaufmann^{1,2}, M. A. Hutterli⁶, T. F. Stocker^{1,2}, U. Ruth⁷, J. P. Steffensen⁵ & V. Maggi³




Figure 1 | EDC dust data in comparison with other climatic indicators. **a**, Stable isotope (δD) record from the EPICA Dome C (EDC) ice core⁸ back to Marine Isotopic Stage 20 (EDC3 timescale) showing Quaternary temperature variations in Antarctica. **b**, Vostok dust flux record (Coulter counter) plotted on its original timescale¹¹. **c**, EDC dust flux records. Red and grey lines represent, respectively, Coulter counter (55-cm to 6-m resolution) and laser-scattering data (55-cm mean). Numbers indicate

Marine Isotopic Stages. Note that the vertical extent of the scales of **b** and **c** is larger than for the other records. **d**, EDC dust size data expressed as FPP (see Methods). The orange and grey curves represent measurements by Coulter counter (2-kyr mean) and laser (1-kyr mean), respectively. **e**, Marine sediment $\delta^{18}O$ stack¹⁸, giving the pattern of global ice volume. **f**, Magnetic susceptibility stack record for Chinese loess¹⁷ (normalized).

Dust depositions on glacial-interglacial time scales




Fig. 2. Iron content fluctuations across the Pacific and Atlantic SO (7) compared to dust content changes in the EDC ice core (1).

SCIENCE VOL 343 24 JANUARY 2014

Increased Dust Deposition in the Pacific Southern Ocean During Glacial Periods

F. Lamy,^{1,2*} R. Gersonde,^{1,2} G. Winckler,^{3,4} O. Esper,¹ A. Jaeschke,^{1,2} G. Kuhn,¹ J. Ullermann,¹ A. Martinez-Garcia,⁵ F. Lambert,⁶ R. Kilian⁷





Fig. 1. Map showing the modern relative contributions of the three major dust sources in the Southern Hemisphere (blue, Australia; red, South America; green, South Africa), based on model data (20). Red dots mark primary core locations; yellow dots indicate additional cores; gray dots denote location of published reference records (1, 4, 7).

LGM dust cycle: fertilisation of the marine biosphere



John Martin

The Iron Hypothesis





Figure 1. In the equatorial Pacific Ocean and Gulf of Alaska, phytoplankton populations are relatively low (purple shaded areas on map), despite adequate sunlight and nutrients. John Martin set out to prove that a lack of dissolved iron in the water in these areas keeps populations of marine algae lower than normal.

- **Dust contains iron which is a key micro-nutrient in the Southern Ocean**
=> higher dust inputs into ocean can fertilise marine biosphere
=> an increased marine bioproductivity will lead to less atmospheric CO₂

LGM dust cycle: fertilisation of the marine biosphere

Ocean Carbon Cycle: Sensitivity Studies for different LGM conditions

- Hypothesis of Martin (1988) : Iron Fertilization of the Glacial Ocean




- increased LGM iron deposition:
-15 ppm


- mechanisms
 - export of organic material
 - export of CaCO₃
(change of alkalinity)

- sensitivity
 - no dust deposition: +3 ppm
 - iron saturation: -39 ppm


Dust in the climate system



Dust in the climate system





Dust in the climate system



LGM climate of North America and Europe



A Modern vegetation




B Glacial vegetation

remember
LGM lecture!

Simulation of glacial dust emissions


modern mean dust emission [$\text{g}/\text{m}^2/\text{yr}$]



changes of glacial dust emissions:

- global emission increase of factor $\sim 2-3$
- glacial increase is not uniform, but varies for the different source regions

LGM mean dust emission [$\text{g}/\text{m}^2/\text{yr}$]



- **Sahara:** increased dust emissions by increased number and intensity of dust storms
- **Asia:** increased dust emissions by increase of (potential) source areas
- seasonal changes of LGM dust emissions are mainly related to changes in vegetation cover

The glacial dust cycle: summary

- **during LGM: 3- to 5-fold increase of dust cycle intensity (global value)**
 - about 1/2 - 2/3 of glacial increase in dust emissions might be caused by the increased wind strength during the LGM
 - about 1/3 - 1/2 of the glacial dust emissions might stem from glacial-only source regions (change of glacial vegetation cover)
- **mineral dust aerosol might be responsible for about 1/4 of the glacial temperature cooling in the (sub)tropical low latitudes**
 - in higher northern latitudes these radiative effects are minor as compared to radiative changes caused by the glacial ice sheet
- **model simulations with marine biogeochemistry models reveal that the glacial atmospheric CO₂ concentration may have been decreased by up to 40-50ppm**
 - it seems unlikely that glacial dust input into the oceans is the only reason for the observed total glacial-interglacial CO₂-reduction of ~80-100 ppm

Climate System II

(Winter 2023/2024)

6th lecture:

Biogeochemical cycles, vegetation and dust

(Aridity and dust, vegetation dynamics, land use, terrestrial biosphere)

End of lecture.

Slides available at:

https://paleodyn.uni-bremen.de/study/climate2023_24.html